

Table 1C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Demographic Indicators
Children Aged < 5 Years (2,3)

Page: 1
Run Date: 3 / 13 / 09

| Total Records | Number |
|----------------------------|---------|
| Individual Children | 129,648 |
| Total Records (All Visits) | 224,336 |

| Source of Data | Number | % |
|----------------------------------|----------------|--------------|
| WIC | 129,648 | 100.0 |
| EPSDT | 0 | 0.0 |
| Child Health-MCH | 0 | 0.0 |
| Child Health-HMO | 0 | 0.0 |
| Headstart | 0 | 0.0 |
| School Health | 0 | 0.0 |
| Other | 0 | 0.0 |
| Total | 129,648 | 100.0 |
| <i>Excluded Unknown:</i> 0 0.0 % | | |

| Race/Ethnic Distribution | Number | % |
|---------------------------------|----------------|--------------|
| White, Not Hispanic | 58,518 | 45.2 |
| Black, Not Hispanic | 24,038 | 18.5 |
| Hispanic | 32,318 | 24.9 |
| American Indian/Alaskan Native | 2,259 | 1.7 |
| Asian/Pacific Islander | 5,674 | 4.4 |
| Multiple Races | 6,798 | 5.2 |
| All Other | 0 | 0.0 |
| Total | 129,605 | 100.0 |
| <i>Excluded Unknown:</i> 43 * % | | |

| Age Distribution (4) | Number | % |
|----------------------|----------------|--------------|
| 0 - 5 Months | 33,598 | 25.9 |
| 6 - 11 Months | 11,842 | 9.1 |
| 12 - 23 Months | 27,481 | 21.2 |
| 24 - 35 Months | 21,327 | 16.4 |
| 3 - 4 Years | 35,400 | 27.3 |
| Total | 129,648 | 100.0 |

| % of Poverty Level | Number | % |
|--------------------------------------|----------------|--------------|
| 0-50 | 27,764 | 21.6 |
| 51-100 | 21,074 | 16.4 |
| 101-130 | 11,185 | 8.7 |
| 131-150 | 6,192 | 4.8 |
| 151-185 | 7,935 | 6.2 |
| 186-200 | 0 | 0.0 |
| Over 200 | 0 | 0.0 |
| Adjunctive Eligibility | 54,458 | 42.3 |
| Total | 128,608 | 100.0 |
| <i>Excluded Unknown:</i> 1,040 0.8 % | | |

| Program Participation | Number | % |
|--------------------------|---------|-------|
| WIC | 129,648 | 100.0 |
| FoodStamps | 123,532 | 43.6 |
| Medicaid | 128,297 | 47.0 |
| TANF | 128,361 | 1.2 |
| <i>Excluded Unknown:</i> | | |
| WIC | 0 | 0.0 |
| FoodStamps | 6,116 | 4.7 |
| Medicaid | 1,351 | 1.0 |
| TANF | 1,287 | 1.0 |

| Migrant Status | Number | % |
|----------------------------------|----------------|--------------|
| Yes | 144 | 0.1 |
| No | 129,504 | 99.9 |
| Total | 129,648 | 100.0 |
| <i>Excluded Unknown:</i> 0 0.0 % | | |

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 2C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Health Indicators
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Summary of Health Indicators ⁽³⁾ | | | | | |
|--|---------------|--|-------------------------|------|--|
| | | | Wisconsin Prevalence | | National Prevalence (Prior Year) |
| | | | Number | % | Number % |
| Birthweight (4) | | | | | |
| Low | < 2500 g | | 31,940 | 8.5 | 1,953,242 9.1 |
| High | > 4000 g | | 31,940 | 7.6 | 1,953,242 6.4 |
| Height and Weight | | | | | |
| Short Stature (5) | < 5th | | 126,769 | 5.5 | 7,526,585 6.2 |
| | < 10th | | 126,769 | 10.5 | 7,526,585 11.2 |
| Underweight (6) | < 5th | | 126,769 | 3.5 | 7,526,585 4.5 |
| | < 10th | | 126,769 | 6.2 | 7,526,585 7.8 |
| Obese (6) ≥ 2 Yrs Overweight (6) | ≥ 95th | | 126,769 | 13.3 | 7,526,585 13.9 |
| | 85th - < 95th | | 55,875 | 16.3 | 3,058,130 16.4 |
| | ≥ 95th | | 55,875 | 13.6 | 3,058,130 14.9 |
| Anemia (7) | | | | | |
| Low Hb | | | 87,908 | 11.3 | 4,357,486 15.3 |
| Low Hct | | | 4,140 | 4.4 | 627,528 11.0 |
| Low Hb/Hct | | | 91,619 | 11.0 | 4,759,588 15.0 |
| Breastfeeding (8) | | | | | |
| Ever Breastfed | | | 31,985 | 64.6 | 1,747,004 59.8 |
| Breastfed At Least 6 Months | | | 13,963 | 27.1 | 874,434 25.4 |
| Breastfed At Least 12 Months | | | 19,862 | 12.3 | 979,105 17.5 |
| Smoking in Household | | | | | |
| ≤ 2 Hours TV Viewing/ Day (9) | | | 127,950 | 9.3 | 3,047,308 11.3 |
| | | | 60,419 | 81.8 | 687,613 75.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(9) Children aged 2 years and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 3C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Breastfeeding Indicators
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Summary of Breastfeeding Indicators ⁽³⁾ | | | | |
|---|-------------------------|------|--|------|
| | Wisconsin Prevalence | | National Prevalence (Prior Year) | |
| | Number | % | Number | % |
| Breastfeeding Duration ^(4,5) | | | | |
| Ever Breastfed | 31,985 | 64.6 | 1,747,004 | 59.8 |
| Breastfed 1+ Weeks | 28,947 | 62.0 | 1,441,655 | 57.7 |
| Breastfed 2+ Weeks | 24,626 | 62.3 | 1,143,026 | 54.6 |
| Breastfed 3+ Weeks | 22,205 | 59.9 | 945,918 | 51.9 |
| Breastfed 4+ Weeks | 20,584 | 57.1 | 829,522 | 49.3 |
| Breastfed 6+ Weeks | 15,634 | 55.3 | 703,467 | 43.4 |
| Breastfed 2+ Months | 12,493 | 54.2 | 656,316 | 38.1 |
| Breastfed 3+ Months | 11,034 | 48.7 | 698,226 | 33.2 |
| Breastfed 4+ Months | 11,477 | 40.3 | 781,546 | 29.4 |
| Breastfed 5+ Months | 12,637 | 32.3 | 876,545 | 26.5 |
| Breastfed 6+ Months | 13,963 | 27.1 | 874,434 | 25.4 |
| Breastfed 9+ Months | 18,210 | 17.5 | 956,555 | 20.2 |
| Breastfed 12+ Months | 19,862 | 12.3 | 979,105 | 17.5 |
| Breastfed 18+ Months | 15,377 | 6.7 | 722,829 | 10.5 |
| Exclusive Breastfeeding ⁽⁶⁾ | | | | |
| Exclusively Breastfed at Least 3 Months | 10,160 | 15.7 | 127,047 | 10.2 |
| Exclusively Breastfed at Least 6 Months | 12,963 | 6.3 | 182,064 | 5.2 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. For each breastfeeding duration category, the analysis includes only children who turned that age during the reporting period by/on their date of visit. For example, infants who turned 9 months old during the reporting period by/on their date of visit are included in the 9+ months analysis.

(5) It is possible the analysis will show that slightly more infants are breastfed at older ages, e.g. 1+ week of age, than at younger ages, e.g. at birth (ever breastfed), due to variation in infants included in each breastfeeding analysis category.

(6) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Contributor | | | | | | | | |
|--|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Contributor | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| Wisconsin | 129,648 | 45.1 | 18.5 | 24.9 | 1.7 | 4.4 | 5.2 | 0.0 |
| Nation (Prior Year) | 7,995,600 | 31.6 | 19.4 | 40.9 | 1.0 | 2.5 | 1.8 | 2.7 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Contributor
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Contributor | | | | |
|--|--|------------------------|-------------------------|-------------------------|
| Contributor | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Wisconsin | 129,648 | 35.0 | 21.2 | 43.8 |
| Nation (Prior Year) | 7,995,600 | 37.6 | 22.1 | 40.3 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Contributor ⁽³⁾ | | | | | | | | | | | | |
|--|----------------------------|--------------|--------------|------------------------------------|-----------|------------------------------------|-------------------------|---|---------------------|------------|--|------|
| Contributor | Birthweight ⁽⁴⁾ | | | Height and Weight | | | | | | | Anemia Low Hb/Hct ⁽⁷⁾ | |
| | Low | | High | Short Stature ⁽⁵⁾ | | Under- weight ⁽⁶⁾ | Obese ⁽⁶⁾ | ≥ 2 Yrs Over- weight ⁽⁶⁾ | | | | |
| | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th- <95th | % ≥95th | Number | % |
| Wisconsin | 31,940 | 8.5 | 7.6 | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 | 11.0 |
| Nation (Prior Year) | 1,953,242 | 9.1 | 6.4 | 7,526,585 | 6.2 | 4.5 | 13.9 | 3,058,130 | 16.4 | 14.9 | 4,759,588 | 15.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children..

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor
(4)

| Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Contributor ⁽⁴⁾ | | | | | | | | | | | | | | |
|---|------------------------------|------------------|---------|-------------------------------|---------|--------------------------------|--|-------------------------------|---------|-------------------------------|--|------|----------------------|------|
| | Breastfeeding ⁽⁵⁾ | | | | | | Exclusive Breastfeeding ⁽⁶⁾ | | | | ≤2 Hours TV Viewing/Day ⁽⁷⁾ | | Smoking in Household | |
| Contributor | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| Wisconsin | 31,985 | 64.6 | 13,963 | 27.1 | 19,862 | 12.3 | 10,160 | 15.7 | 12,963 | 6.3 | 60,419 | 81.8 | 127,950 | 9.3 |
| Nation (Prior Year) | 1,747,004 | 59.8 | 874,434 | 25.4 | 979,105 | 17.5 | 127,047 | 10.2 | 182,064 | 5.2 | 687,613 | 75.0 | 3,047,308 | 11.3 |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(6) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 8C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Growth and Anemia Indicators by Race/Ethnicity or Age
Children Aged < 5 Years ⁽²⁾

| Growth and Anemia Indicators by Race/Ethnicity or Age ⁽³⁾ | | | | | | | | | | | | |
|--|-----------------|----------|-----------|-------------------|-------------------------|-------------------------|--------------|------------------------------|-----------------|------------|-----------------------------|------|
| | Birthweight (4) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | Number | Low % | High % | Number | Short Stature (5) | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | |
| | | | | | | | | | | | | |
| | | <2500 g | >4000 g | | % <5th | % <5th | % ≥95th | Number | % 85th-<95th | % ≥95th | Number | % |
| Race/Ethnicity | | | | | | | | | | | | |
| White, Not Hispanic | 15,021 | 8.0 | 9.0 | 57,393 | 5.3 | 3.6 | 11.5 | 24,400 | 16.0 | 11.3 | 40,498 | 7.6 |
| Black, Not Hispanic | 6,211 | 12.1 | 3.9 | 23,410 | 6.0 | 4.9 | 11.4 | 10,078 | 13.5 | 10.2 | 16,586 | 22.8 |
| Hispanic | 7,335 | 6.5 | 8.2 | 31,576 | 4.9 | 2.4 | 16.8 | 14,691 | 18.0 | 18.5 | 23,758 | 9.3 |
| American Indian/Alaskan Native | 423 | 5.2 | 12.3 | 2,188 | 3.2 | 2.6 | 21.0 | 1,103 | 21.0 | 24.1 | 1,738 | 7.4 |
| Asian/Pacific Islander | 1,337 | 7.6 | 3.3 | 5,549 | 10.4 | 2.8 | 14.5 | 2,595 | 17.2 | 16.2 | 4,119 | 9.2 |
| Multiple Races | 1,613 | 9.1 | 8.4 | 6,639 | 4.5 | 3.3 | 14.6 | 2,994 | 16.8 | 13.0 | 4,895 | 10.2 |
| All Other | 0 | * | * | 14 | * | * | * | 14 | * | * | 25 | * |
| Total | 31,940 | 8.5 | 7.6 | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 | 11.0 |
| Age | | | | | | | | | | | | |
| 0 - 11 Months | - | - | - | 43,214 | 7.5 | 4.9 | 10.5 | - | * | * | 5,410 | 16.2 |
| 12 - 23 Months | - | - | - | 27,680 | 5.6 | 2.0 | 16.9 | - | * | * | 32,563 | 14.9 |
| 24 - 35 Months | - | - | - | 21,266 | 4.2 | 2.9 | 12.2 | 21,266 | 15.7 | 12.2 | 21,905 | 10.2 |
| 36 - 47 Months | - | - | - | 17,663 | 3.9 | 3.2 | 13.8 | 17,663 | 16.5 | 13.8 | 17,883 | 7.2 |
| 48 - 59 Months | - | - | - | 16,946 | 3.6 | 3.0 | 15.1 | 16,946 | 16.7 | 15.1 | 13,858 | 5.9 |
| Total | - | - | - | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 | 11.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 9C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (2)
Children Aged < 5 Years (3)

| Breastfeeding, TV Viewing, and Smoking in Household by Race/Ethnicity or Age (4) | | | | | | | | | | |
|--|-------------------|------------------|--------|-------------------------------|--------|--------------------------------|-----------------------------------|------|----------------------|------|
| | Breastfeeding (5) | | | | | | <2 Hours TV/Video Viewing/Day (6) | | Smoking in Household | |
| | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % | Number | % |
| Race/Ethnicity | | | | | | | | | | |
| White, Not Hispanic | 15,057 | 66.0 | 6,543 | 27.0 | 8,782 | 11.2 | 26,253 | 84.6 | 57,886 | 10.0 |
| Black, Not Hispanic | 6,222 | 50.4 | 2,447 | 13.4 | 3,810 | 5.8 | 10,811 | 76.1 | 23,615 | 14.8 |
| Hispanic | 7,334 | 78.6 | 3,503 | 40.2 | 5,002 | 21.0 | 16,115 | 81.7 | 31,913 | 4.0 |
| American Indian/Alaskan Native | 425 | 58.8 | 221 | 21.3 | 316 | 11.7 | 1,196 | 82.4 | 2,222 | 12.3 |
| Asian/Pacific Islander | 1,338 | 44.6 | 546 | 16.8 | 876 | 7.4 | 2,803 | 77.1 | 5,614 | 2.8 |
| Multiple Races | 1,609 | 60.0 | 703 | 20.9 | 1,076 | 8.1 | 3,233 | 82.0 | 6,692 | 12.3 |
| All Other | 0 | * | 0 | * | 0 | * | 8 | * | 8 | * |
| Total | 31,985 | 64.6 | 13,963 | 27.1 | 19,862 | 12.3 | 60,419 | 81.8 | 127,950 | 9.3 |
| Age | | | | | | | | | | |
| 0 - 11 Months | - | - | - | - | - | - | 0 | * | 43,904 | 7.4 |
| 12 - 23 Months | - | - | - | - | - | - | 0 | * | 27,430 | 9.3 |
| 24 - 35 Months | - | - | - | - | - | - | 25,184 | 85.3 | 21,287 | 10.3 |
| 36 - 47 Months | - | - | - | - | - | - | 17,928 | 79.3 | 17,992 | 11.1 |
| 48 - 59 Months | - | - | - | - | - | - | 17,307 | 79.2 | 17,337 | 10.6 |
| Total | 31,985 | 64.6 | 13,963 | 27.1 | 19,862 | 12.3 | 60,419 | 81.8 | 127,950 | 9.3 |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 10C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Summary of Trends in Racial and Ethnic Distribution | | | | | | | | |
|--|---------------------------------------|-----------------------------|-----------------------------|---------------|---|--------------------------------|---------------------|-----------------------------|
| Year | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| 2008 | 129,648 | 45.1 | 18.5 | 24.9 | 1.7 | 4.4 | 5.2 | 0.0 |
| 2007 | 122,799 | 45.2 | 18.6 | 24.7 | 1.9 | 4.4 | 5.1 | 0.0 |
| 2006 | 121,673 | 45.1 | 18.6 | 24.2 | 2.0 | 4.5 | 5.1 | 0.5 |
| 2005 | 115,967 | 49.6 | 20.3 | 20.7 | 2.4 | 4.7 | 1.0 | 1.3 |
| 2004 | 116,740 | 52.2 | 21.5 | 19.2 | 2.5 | 4.6 | 0.0 | 0.0 |
| 2003 | 112,855 | 52.8 | 22.0 | 18.3 | 2.5 | 4.4 | 0.0 | 0.0 |
| 2002 | 110,533 | 53.1 | 22.3 | 17.5 | 2.5 | 4.6 | 0.0 | 0.0 |
| 2001 | 108,998 | 53.1 | 23.1 | 16.4 | 2.5 | 4.9 | 0.0 | 0.0 |
| 2000 | 106,063 | 53.7 | 23.7 | 14.8 | 2.5 | 5.3 | 0.0 | 0.0 |
| 1999 | 104,221 | 54.4 | 24.5 | 13.0 | 2.5 | 5.7 | 0.0 | 0.0 |
| 1998 | 111,982 | 55.0 | 24.8 | 11.6 | 2.5 | 6.0 | 0.0 | 0.0 |
| 1997 | 116,090 | 55.2 | 25.2 | 10.7 | 2.6 | 6.4 | 0.0 | 0.0 |
| 1996 | 117,936 | 54.9 | 25.6 | 10.2 | 2.5 | 6.8 | 0.0 | 0.0 |
| 1995 | 119,022 | 54.6 | 26.4 | 9.5 | 2.4 | 7.2 | 0.0 | 0.0 |
| 1994 | 119,132 | 55.4 | 26.5 | 8.4 | 2.4 | 7.3 | 0.0 | 0.0 |
| 1993 | 272 | 46.7 | 33.1 | 14.0 | 1.1 | 5.1 | 0.0 | 0.0 |
| 1992 | 0 | * | * | * | * | * | * | * |
| 1991 | 0 | * | * | * | * | * | * | * |
| 1990 | 0 | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 10C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Racial and Ethnic Distribution
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3 / 13 / 09

| Summary of Trends in Racial and Ethnic Distribution | | | | | | | | |
|--|---------------------------------------|-----------------------------|-----------------------------|---------------|---|--------------------------------|---------------------|-----------------------------|
| Year | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| 1989 | 0 | * | * | * | * | * | * | * |
| 1988 | 0 | * | * | * | * | * | * | * |
| 1987 | 0 | * | * | * | * | * | * | * |
| 1986 | 0 | * | * | * | * | * | * | * |
| 1985 | 0 | * | * | * | * | * | * | * |
| 1984 | 0 | * | * | * | * | * | * | * |
| 1983 | 0 | * | * | * | * | * | * | * |
| 1982 | 115 | 69.6 | 22.6 | 0.0 | 2.6 | 5.2 | 0.0 | 0.0 |
| 1981 | 14,903 | 62.4 | 16.8 | 6.9 | 10.4 | 3.3 | 0.0 | 0.0 |
| 1980 | 31,119 | 68.5 | 13.8 | 6.6 | 8.7 | 2.4 | 0.0 | 0.1 |
| 1979 | 22,015 | 64.0 | 17.8 | 7.0 | 9.4 | 1.5 | 0.0 | 0.3 |
| 1978 | 377 | 97.9 | 0.3 | 0.3 | 0.3 | 0.5 | 0.0 | 0.8 |
| 1977 | 9 | * | * | * | * | * | * | * |
| 1976 | 5 | * | * | * | * | * | * | * |
| 1975 | 3 | * | * | * | * | * | * | * |
| 1974 | 1 | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Summary of Trends in Age Distribution (3) | | | | |
|--|--|------------------|-------------------|-------------------|
| Year | Number Included in Analysis (4) | 0-11 Months % | 12-23 Months % | 24-59 Months % |
| 2008 | 129,648 | 35.0 | 21.2 | 43.8 |
| 2007 | 122,799 | 36.3 | 21.1 | 42.6 |
| 2006 | 121,673 | 36.7 | 19.9 | 43.3 |
| 2005 | 115,967 | 31.6 | 23.2 | 45.2 |
| 2004 | 116,740 | 31.5 | 23.1 | 45.4 |
| 2003 | 112,855 | 31.8 | 22.7 | 45.5 |
| 2002 | 110,533 | 31.6 | 23.0 | 45.4 |
| 2001 | 108,998 | 32.2 | 23.0 | 44.8 |
| 2000 | 106,063 | 31.5 | 23.3 | 45.2 |
| 1999 | 104,221 | 30.2 | 23.1 | 46.7 |
| 1998 | 111,982 | 30.2 | 22.4 | 47.4 |
| 1997 | 116,090 | 29.1 | 22.1 | 48.8 |
| 1996 | 117,936 | 28.4 | 22.4 | 49.2 |
| 1995 | 119,022 | 28.7 | 22.2 | 49.1 |
| 1994 | 119,132 | 33.2 | 19.6 | 47.2 |
| 1993 | 272 | 86.8 | 6.3 | 7.0 |
| 1992 | 0 | * | * | * |
| 1991 | 0 | * | * | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Records rejected if date of birth and date of visit unknown.

(4) Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 11C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Age Distribution
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 13 / 09

| Summary of Trends in Age Distribution (3) | | | | |
|--|--|------------------|-------------------|-------------------|
| Year | Number Included in Analysis (4) | 0-11 Months % | 12-23 Months % | 24-59 Months % |
| 1990 | 0 | * | * | * |
| 1989 | 0 | * | * | * |
| 1988 | 0 | * | * | * |
| 1987 | 0 | * | * | * |
| 1986 | 0 | * | * | * |
| 1985 | 0 | * | * | * |
| 1984 | 0 | * | * | * |
| 1983 | 0 | * | * | * |
| 1982 | 115 | 79.1 | 5.2 | 15.7 |
| 1981 | 14,903 | 43.5 | 26.9 | 29.5 |
| 1980 | 31,119 | 34.4 | 21.6 | 44.0 |
| 1979 | 22,015 | 29.7 | 21.6 | 48.7 |
| 1978 | 377 | 1.9 | 2.1 | 96.0 |
| 1977 | 9 | * | * | * |
| 1976 | 5 | * | * | * |
| 1975 | 3 | * | * | * |
| 1974 | 1 | * | * | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Records rejected if date of birth and date of visit unknown.

(4) Excludes records with unknown data or errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 12C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3/13/2009

| Summary of Trends in Growth and Anemia Indicators ⁽³⁾ | | | | | | | | | | | | |
|--|-----------------|--------------|--------------|-------------------------|-----------|-------------------------|--------------|------------------------------|-----------------|------------|-----------------------------|------|
| Year | Birthweight (4) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | Low | | High | Short Stature (5) | | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | |
| | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th-<95th | % ≥95th | Number | % |
| 2008 | 31,940 | 8.5 | 7.6 | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 | 11.0 |
| 2007 | 31,162 | 8.7 | 7.5 | 120,071 | 5.4 | 3.7 | 12.2 | 51,636 | 16.1 | 13.1 | 84,006 | 11.3 |
| 2006 | 31,039 | 8.6 | 7.6 | 118,548 | 5.7 | 3.8 | 12.4 | 51,667 | 16.3 | 13.0 | 85,045 | 10.4 |
| 2005 | 28,451 | 8.6 | 8.1 | 112,753 | 5.5 | 4.5 | 12.3 | 51,410 | 16.0 | 13.0 | 77,037 | 12.2 |
| 2004 | 29,547 | 8.7 | 8.3 | 113,275 | 5.4 | 4.3 | 12.4 | 51,825 | 16.3 | 13.3 | 78,255 | 13.2 |
| 2003 | 28,471 | 8.3 | 8.5 | 109,384 | 5.2 | 4.4 | 12.2 | 50,284 | 15.9 | 13.0 | 75,448 | 12.2 |
| 2002 | 27,201 | 8.5 | 8.7 | 106,429 | 5.5 | 4.3 | 11.8 | 48,993 | 15.0 | 11.8 | 74,420 | 12.7 |
| 2001 | 27,782 | 8.4 | 8.9 | 104,796 | 5.6 | 4.6 | 11.4 | 47,741 | 14.6 | 11.4 | 71,952 | 13.0 |
| 2000 | 27,000 | 8.5 | 8.9 | 102,004 | 5.5 | 4.5 | 11.2 | 46,744 | 15.1 | 11.5 | 70,955 | 13.6 |
| 1999 | 25,565 | 8.8 | 9.0 | 100,387 | 5.4 | 4.4 | 10.5 | 47,584 | 14.6 | 10.1 | 69,517 | 13.5 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3/13/2009

| Summary of Trends in Growth and Anemia Indicators ⁽³⁾ | | | | | | | | | | | | |
|--|-----------------|--------------|--------------|-------------------------|-------------------------|--------------|------------------------------|--------|-----------------|------------|-----------------------------|------|
| Year | Birthweight (4) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | Low | High | | Short Stature (5) | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | | |
| | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th-<95th | % ≥95th | Number | % |
| 1998 | 27,014 | 8.6 | 9.1 | 109,030 | 5.6 | 4.5 | 10.2 | 52,186 | 14.4 | 10.1 | 77,025 | 13.3 |
| 1997 | 26,681 | 8.5 | 9.2 | 113,120 | 5.6 | 4.3 | 10.2 | 55,828 | 14.7 | 10.1 | 82,877 | 13.1 |
| 1996 | 26,542 | 8.3 | 9.4 | 114,905 | 5.8 | 4.6 | 9.6 | 57,370 | 14.6 | 9.7 | 85,637 | 14.0 |
| 1995 | 26,268 | 8.4 | 9.2 | 116,002 | 6.3 | 4.8 | 9.4 | 57,724 | 13.9 | 9.3 | 87,361 | 16.2 |
| 1994 | 26,120 | 8.8 | 9.4 | 116,138 | 6.5 | 4.9 | 9.4 | 55,382 | 13.8 | 8.9 | 88,300 | 17.2 |
| 1993 | 231 | 10.0 | 8.2 | 253 | 1.2 | 11.9 | 5.5 | 19 | * | * | 45 | * |
| 1992 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1991 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1990 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1989 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3/13/2009

| Summary of Trends in Growth and Anemia Indicators ⁽³⁾ | | | | | | | | | | | | |
|--|-----------------|--------------|--------------|-------------------------|-----------|-------------------------|--------------|------------------------------|-----------------|-----------------------------|--------|------|
| Year | Birthweight (4) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (7) | | |
| | Low | | High | Short Stature (5) | | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | |
| | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th-<95th | % ≥95th | Number | % |
| 1988 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1987 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1986 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1985 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1984 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1983 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| 1982 | 69 | * | * | 101 | 5.9 | 10.9 | 4.0 | 17 | * | * | 33 | * |
| 1981 | 3,137 | 7.8 | 11.9 | 12,712 | 7.5 | 4.9 | 9.7 | 4,276 | 14.0 | 7.6 | 10,575 | 10.8 |
| 1980 | 6,007 | 7.7 | 11.8 | 27,903 | 6.5 | 5.0 | 9.0 | 13,338 | 13.4 | 7.6 | 23,768 | 8.3 |
| 1979 | 4,252 | 8.5 | 10.8 | 19,769 | 6.2 | 4.8 | 10.1 | 10,382 | 13.3 | 8.4 | 16,871 | 9.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3/13/2009

| Summary of Trends in Growth and Anemia Indicators ⁽³⁾ | | | | | | | | | | | | |
|--|-----------------|--------------|--------------|-------------------------|-----------|-------------------------|--------------|------------------------------|-----------------|-----------------------------|--------|---|
| Year | Birthweight (4) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (7) | | |
| | Low | | High | Short Stature (5) | | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | |
| | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th-<95th | % ≥95th | Number | % |
| 1978 | 2 | * | * | 366 | 3.6 | 4.1 | 8.5 | 354 | 17.5 | 8.5 | 29 | * |
| 1977 | 2 | * | * | 0 | * | * | * | 0 | * | * | 5 | * |
| 1976 | 4 | * | * | 0 | * | * | * | 0 | * | * | 1 | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 13C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Breastfeeding
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3/13/2009

| Summary of Trends in Breastfeeding ⁽³⁾ | | | | | | |
|--|----------------------|------------------------|--------|--|--------|---|
| Year | Breastfeeding (4) | | | | | |
| | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| 2008 | 31,985 | 64.6 | 13,963 | 27.1 | 19,862 | 12.3 |
| 2007 | 31,171 | 63.2 | 13,616 | 26.6 | 18,952 | 12.3 |
| 2006 | 31,071 | 62.6 | 12,726 | 26.0 | 17,467 | 13.2 |
| 2005 | 28,610 | 59.9 | 10,409 | 24.2 | 17,417 | 16.4 |
| 2004 | 27,213 | 59.3 | 11,238 | 25.0 | 17,605 | 16.1 |
| 2003 | 28,288 | 58.2 | 10,556 | 23.8 | 16,675 | 14.8 |
| 2002 | 27,000 | 55.0 | 10,711 | 22.6 | 16,754 | 14.9 |
| 2001 | 27,630 | 53.4 | 10,314 | 22.5 | 16,396 | 14.4 |
| 2000 | 26,838 | 51.0 | 9,868 | 22.1 | 16,194 | 14.6 |
| 1999 | 25,360 | 49.9 | 9,857 | 20.0 | 15,919 | 10.9 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 13C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Breastfeeding
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3/13/2009

| Summary of Trends in Breastfeeding ⁽³⁾ | | | | | | |
|--|----------------------|------------------------|--------|--|--------|---|
| Year | Breastfeeding (4) | | | | | |
| | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| 1998 | 26,682 | 50.4 | 10,148 | 17.9 | 16,018 | 7.9 |
| 1997 | 26,338 | 48.7 | 10,663 | 17.3 | 16,548 | 9.8 |
| 1996 | 25,997 | 48.7 | 10,693 | 17.4 | 16,640 | 10.4 |
| 1995 | 25,753 | 47.0 | 10,973 | 18.6 | 16,161 | 12.2 |
| 1994 | 25,798 | 44.8 | 13,675 | 14.0 | 14,031 | 8.4 |
| 1993 | 226 | 42.0 | 14 | * | 3 | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 14C
2008 Pediatric Nutrition Surveillance
Wisconsin
Children Aged < 5 Years

Page: 1
Run Date: 3 / 13 / 09

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 15C
2008 Pediatric Nutrition Surveillance
Wisconsin
Children Aged < 5 Years

Page: 1
Run Date: 3 / 13 / 09

This table is forthcoming. It may present changes over time in the prevalence of short stature, underweight, overweight, or anemia among cohorts of children participating in the PedNSS.

Table 16C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3/13/2009

| Growth Indicators by Race/Ethnicity and Age ⁽³⁾ | | | | | | | | |
|---|----------------------|------------|--------------------|------------|---------------|-------------|---------------------------|---------------------------------|
| | Short Stature (4) | | Underweight (5) | | Obese (5) | | ≥ 2 Yrs Overweight (5) | |
| | Number | % <5th | Number | % <5th | Number | % ≥95th | Number | % 85th - <95th % ≥95th |
| White, Not Hispanic | | | | | | | | |
| 0 - 11 Months | 20,785 | 7.2 | 20,785 | 4.8 | 20,785 | 9.5 | 0 | 0.0 0.0 |
| 12 - 23 Months | 12,208 | 5.3 | 12,208 | 2.0 | 12,208 | 15.4 | 0 | 0.0 0.0 |
| 24 - 59 Months | 24,400 | 3.8 | 24,400 | 3.3 | 24,400 | 11.3 | 24,400 | 16.0 11.3 |
| Total | 57,393 | 5.3 | 57,393 | 3.6 | 57,393 | 11.5 | 24,400 | 16.0 11.3 |
| Black, Not Hispanic | | | | | | | | |
| 0 - 11 Months | 8,051 | 9.8 | 8,051 | 6.6 | 8,051 | 9.8 | 0 | 0.0 0.0 |
| 12 - 23 Months | 5,281 | 6.1 | 5,281 | 2.8 | 5,281 | 16.0 | 0 | 0.0 0.0 |
| 24 - 59 Months | 10,078 | 2.9 | 10,078 | 4.6 | 10,078 | 10.2 | 10,078 | 13.5 10.2 |
| Total | 23,410 | 6.0 | 23,410 | 4.9 | 23,410 | 11.4 | 10,078 | 13.5 10.2 |
| Hispanic | | | | | | | | |
| 0 - 11 Months | 9,949 | 6.2 | 9,949 | 3.9 | 9,949 | 12.2 | 0 | 0.0 0.0 |
| 12 - 23 Months | 6,936 | 4.9 | 6,936 | 1.4 | 6,936 | 19.6 | 0 | 0.0 0.0 |
| 24 - 59 Months | 14,691 | 4.1 | 14,691 | 1.9 | 14,691 | 18.5 | 14,691 | 18.0 18.5 |
| Total | 31,576 | 4.9 | 31,576 | 2.4 | 31,576 | 16.8 | 14,691 | 18.0 18.5 |
| American Indian/Alaskan Native | | | | | | | | |
| 0 - 11 Months | 583 | 5.3 | 583 | 6.5 | 583 | 12.3 | 0 | 0.0 0.0 |
| 12 - 23 Months | 502 | 3.2 | 502 | 1.8 | 502 | 24.1 | 0 | 0.0 0.0 |
| 24 - 59 Months | 1,103 | 2.2 | 1,103 | 0.8 | 1,103 | 24.1 | 1,103 | 21.0 24.1 |
| Total | 2,188 | 3.2 | 2,188 | 2.6 | 2,188 | 21.0 | 1,103 | 21.0 24.1 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 16C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Growth Indicators by Race/Ethnicity and Age
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3/13/2009

| Growth Indicators by Race/Ethnicity and Age ⁽³⁾ | | | | | | | | |
|---|----------------------|-------------|--------------------|------------|----------------|-------------|---------------------------|---------------------------------|
| | Short Stature (4) | | Underweight (5) | | Obese (5) | | ≥ 2 Yrs Overweight (5) | |
| | Number | % <5th | Number | % <5th | Number | % ≥95th | Number | % 85th - <95th % ≥95th |
| Asian/Pacific Islander | | | | | | | | |
| 0 - 11 Months | 1,720 | 8.6 | 1,720 | 4.1 | 1,720 | 11.0 | 0 | 0.0 0.0 |
| 12 - 23 Months | 1,234 | 12.3 | 1,234 | 2.4 | 1,234 | 15.7 | 0 | 0.0 0.0 |
| 24 - 59 Months | 2,595 | 10.7 | 2,595 | 2.2 | 2,595 | 16.2 | 2,595 | 17.2 16.2 |
| Total | 5,549 | 10.4 | 5,549 | 2.8 | 5,549 | 14.5 | 2,595 | 17.2 16.2 |
| Multiple Races | | | | | | | | |
| 0 - 11 Months | 2,126 | 7.1 | 2,126 | 4.6 | 2,126 | 13.4 | 0 | 0.0 0.0 |
| 12 - 23 Months | 1,519 | 4.9 | 1,519 | 1.8 | 1,519 | 19.2 | 0 | 0.0 0.0 |
| 24 - 59 Months | 2,994 | 2.5 | 2,994 | 3.1 | 2,994 | 13.0 | 2,994 | 16.8 13.0 |
| Total | 6,639 | 4.5 | 6,639 | 3.3 | 6,639 | 14.6 | 2,994 | 16.8 13.0 |
| All Other/Unknown | | | | | | | | |
| 0 - 11 Months | 0 | * | 0 | * | 0 | * | 0 | * * |
| 12 - 23 Months | 0 | * | 0 | * | 0 | * | 0 | * * |
| 24 - 59 Months | 14 | * | 14 | * | 14 | * | 14 | * * |
| Total | 14 | * | 14 | * | 14 | * | 14 | * * |
| Total All Race/Ethnic Groups | | | | | | | | |
| 0 - 11 Months | 43,214 | 7.5 | 43,214 | 4.9 | 43,214 | 10.5 | 0 | 0.0 0.0 |
| 12 - 23 Months | 27,680 | 5.6 | 27,680 | 2.0 | 27,680 | 16.9 | 0 | 0.0 0.0 |
| 24 - 59 Months | 55,875 | 3.9 | 55,875 | 3.0 | 55,875 | 13.6 | 55,875 | 16.3 13.6 |
| Total | 126,769 | 5.5 | 126,769 | 3.5 | 126,769 | 13.3 | 55,875 | 16.3 13.6 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 17C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Anemia Indicators by Race/Ethnicity and Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | |
|---|---------------------------|-------------|------------------------------|---------------------------|-------------|
| | Anemia Low Hb/Hct % | | | Anemia Low Hb/Hct % | |
| | Number | | | Number | |
| White, Not Hispanic | | | Asian/Pacific Islander | | |
| 6 - 11 Months | 2,788 | 12.3 | 6 - 11 Months | 234 | 6.8 |
| 12 - 17 Months | 8,146 | 12.4 | 12 - 17 Months | 791 | 11.3 |
| 18 - 23 Months | 6,350 | 8.2 | 18 - 23 Months | 669 | 9.7 |
| 24 - 35 Months | 9,395 | 6.8 | 24 - 35 Months | 960 | 11.9 |
| 36 - 59 Months | 13,819 | 4.0 | 36 - 59 Months | 1,465 | 6.5 |
| Total | 40,498 | 7.6 | Total | 4,119 | 9.2 |
| Black, Not Hispanic | | | Multiple Races | | |
| 6 - 11 Months | 823 | 30.7 | 6 - 11 Months | 264 | 13.3 |
| 12 - 17 Months | 3,441 | 29.3 | 12 - 17 Months | 982 | 17.7 |
| 18 - 23 Months | 2,649 | 26.5 | 18 - 23 Months | 753 | 12.4 |
| 24 - 35 Months | 3,914 | 22.4 | 24 - 35 Months | 1,238 | 9.0 |
| 36 - 59 Months | 5,759 | 16.2 | 36 - 59 Months | 1,658 | 5.0 |
| Total | 16,586 | 22.8 | Total | 4,895 | 10.2 |
| Hispanic | | | All Other/Unknown | | |
| 6 - 11 Months | 1,221 | 17.9 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 4,538 | 16.0 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 3,667 | 10.4 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 5,956 | 8.0 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 8,376 | 4.8 | 36 - 59 Months | 25 | * |
| Total | 23,758 | 9.3 | Total | 25 | * |
| American Indian/Alaskan Native | | | Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 80 | * | 6 - 11 Months | 5,410 | 16.2 |
| 12 - 17 Months | 298 | 14.1 | 12 - 17 Months | 18,196 | 16.8 |
| 18 - 23 Months | 279 | 8.6 | 18 - 23 Months | 14,367 | 12.5 |
| 24 - 35 Months | 442 | 6.1 | 24 - 35 Months | 21,905 | 10.2 |
| 36 - 59 Months | 639 | 3.9 | 36 - 59 Months | 31,741 | 6.6 |
| Total | 1,738 | 7.4 | Total | 91,619 | 11.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

| Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | | | | | | | |
|--|--------|----------------------------|--------------|-------------------------|-------------------------|--------------|------------------------------|------------|--------|-------------|------------|-----------------------------|--------|
| Race/Ethnicity and Year | | Birthweight ⁽⁴⁾ | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | | Low | High | Short Stature (5) | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | | | |
| | | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th - | % <95th | % ≥95th | Number |
| White, Not Hispanic | 2008 | 15,021 | 8.0 | 9.0 | 57,393 | 5.3 | 3.6 | 11.5 | 24,400 | 16.0 | 11.3 | 40,498 | 7.6 |
| | 2007 | 14,548 | 8.1 | 8.6 | 54,396 | 5.4 | 3.8 | 10.8 | 22,789 | 15.7 | 11.0 | 37,129 | 7.5 |
| | 2006 | 14,299 | 8.2 | 8.6 | 53,756 | 5.5 | 3.8 | 10.8 | 23,125 | 15.9 | 10.7 | 38,176 | 6.9 |
| | 2005 | 14,231 | 8.0 | 9.1 | 56,112 | 5.3 | 4.3 | 11.1 | 25,217 | 15.4 | 11.5 | 37,793 | 8.5 |
| | 2004 | 15,915 | 8.5 | 9.4 | 59,280 | 5.3 | 4.1 | 11.3 | 26,229 | 16.0 | 11.9 | 40,143 | 9.3 |
| | 2003 | 15,543 | 7.3 | 9.7 | 57,981 | 5.0 | 4.4 | 11.1 | 26,041 | 15.9 | 11.8 | 39,157 | 8.1 |
| | 2002 | 14,697 | 7.8 | 10.1 | 56,663 | 5.0 | 4.3 | 10.4 | 25,667 | 14.9 | 10.6 | 38,970 | 8.4 |
| | 2001 | 15,042 | 7.8 | 10.2 | 55,794 | 5.2 | 4.5 | 10.3 | 24,906 | 14.0 | 10.4 | 37,577 | 9.5 |
| | 2000 | 14,940 | 7.8 | 10.3 | 54,936 | 5.3 | 4.4 | 10.1 | 24,466 | 14.9 | 10.2 | 37,526 | 9.0 |
| 1999 | 14,391 | 7.6 | 10.6 | 54,699 | 4.9 | 4.3 | 9.6 | 25,110 | 14.5 | 9.2 | 37,247 | 8.2 | |
| Black, Not Hispanic | 2008 | 6,211 | 12.1 | 3.9 | 23,410 | 6.0 | 4.9 | 11.4 | 10,078 | 13.5 | 10.2 | 16,586 | 22.8 |
| | 2007 | 6,214 | 13.4 | 3.9 | 22,250 | 5.7 | 5.1 | 10.3 | 9,433 | 13.3 | 10.2 | 15,245 | 23.7 |
| | 2006 | 6,049 | 12.5 | 4.3 | 22,015 | 6.0 | 5.3 | 10.5 | 9,497 | 14.1 | 10.3 | 15,502 | 21.5 |
| | 2005 | 6,028 | 12.4 | 5.0 | 22,789 | 5.4 | 6.1 | 10.4 | 10,401 | 14.0 | 10.0 | 15,560 | 23.6 |
| | 2004 | 6,485 | 12.3 | 5.3 | 24,278 | 5.6 | 6.1 | 10.6 | 11,241 | 14.3 | 10.7 | 16,818 | 25.4 |
| | 2003 | 6,332 | 12.7 | 5.1 | 23,964 | 5.5 | 5.6 | 10.3 | 11,180 | 13.6 | 10.1 | 16,601 | 22.5 |
| | 2002 | 6,077 | 12.3 | 5.2 | 23,664 | 6.5 | 5.3 | 9.9 | 11,172 | 13.2 | 8.9 | 16,779 | 23.7 |
| | 2001 | 6,302 | 12.0 | 5.4 | 24,070 | 6.4 | 5.6 | 9.8 | 11,281 | 12.9 | 8.2 | 16,980 | 23.0 |
| | 2000 | 6,343 | 12.3 | 5.1 | 24,136 | 5.8 | 5.9 | 9.9 | 11,286 | 13.0 | 8.8 | 16,925 | 25.4 |
| | 1999 | 6,153 | 13.1 | 5.0 | 24,581 | 6.2 | 5.6 | 9.4 | 12,010 | 12.3 | 8.1 | 17,156 | 25.6 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 2
Run Date: 3 / 13 / 09

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

| Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | | | | | | | |
|--|-------|----------------------------|--------------|--------------|-------------------------|-----------|-------------------------|--------------|------------------------------|-------------------|------------|-----------------------------|------|
| Race/Ethnicity and Year | | Birthweight ⁽⁴⁾ | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | | Low | | High | Short Stature (5) | | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | |
| | | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th - <95th | % ≥95th | Number | % |
| Hispanic | 2008 | 7,335 | 6.5 | 8.2 | 31,576 | 4.9 | 2.4 | 16.8 | 14,691 | 18.0 | 18.5 | 23,758 | 9.3 |
| | 2007 | 7,098 | 6.4 | 8.6 | 29,717 | 4.8 | 2.7 | 15.2 | 13,278 | 17.7 | 17.9 | 21,684 | 9.8 |
| | 2006 | 7,226 | 6.7 | 8.8 | 28,571 | 4.9 | 2.7 | 15.6 | 12,876 | 18.5 | 18.1 | 21,121 | 9.1 |
| | 2005 | 5,438 | 6.1 | 9.0 | 23,228 | 4.7 | 3.6 | 16.0 | 11,152 | 18.6 | 17.7 | 16,590 | 10.2 |
| | 2004 | 5,189 | 5.8 | 8.9 | 21,669 | 4.3 | 3.4 | 15.9 | 10,455 | 18.3 | 18.1 | 15,611 | 10.9 |
| | 2003 | 4,844 | 6.3 | 9.0 | 19,959 | 4.7 | 3.5 | 16.0 | 9,407 | 17.6 | 17.8 | 14,379 | 11.5 |
| | 2002 | 4,734 | 6.2 | 9.0 | 18,552 | 4.8 | 3.8 | 16.3 | 8,445 | 16.4 | 16.4 | 13,251 | 11.6 |
| | 2001 | 4,659 | 6.1 | 9.8 | 17,169 | 4.9 | 3.9 | 15.1 | 7,648 | 17.5 | 15.5 | 11,899 | 11.1 |
| | 2000 | 3,975 | 6.5 | 9.8 | 14,962 | 4.6 | 3.7 | 14.3 | 6,812 | 17.6 | 15.9 | 10,611 | 12.5 |
| 1999 | 3,355 | 7.1 | 9.7 | 12,893 | 4.4 | 3.9 | 13.4 | 5,961 | 15.9 | 13.5 | 9,036 | 13.6 | |
| American Indian/Alaskan Native | 2008 | 423 | 5.2 | 12.3 | 2,188 | 3.2 | 2.6 | 21.0 | 1,103 | 21.0 | 24.1 | 1,738 | 7.4 |
| | 2007 | 497 | 4.8 | 11.7 | 2,320 | 3.2 | 2.5 | 19.7 | 1,102 | 22.5 | 21.4 | 1,757 | 6.1 |
| | 2006 | 573 | 6.6 | 12.7 | 2,397 | 3.4 | 2.4 | 20.1 | 1,119 | 21.9 | 21.0 | 1,796 | 7.1 |
| | 2005 | 656 | 6.6 | 16.5 | 2,716 | 3.4 | 2.7 | 18.6 | 1,318 | 21.9 | 19.3 | 1,832 | 7.6 |
| | 2004 | 678 | 4.7 | 14.3 | 2,836 | 3.5 | 3.2 | 19.2 | 1,361 | 22.6 | 20.5 | 1,971 | 8.0 |
| | 2003 | 608 | 5.9 | 17.3 | 2,674 | 4.0 | 2.1 | 19.8 | 1,246 | 20.0 | 19.2 | 1,900 | 8.2 |
| | 2002 | 635 | 4.6 | 13.5 | 2,666 | 3.7 | 2.4 | 18.3 | 1,286 | 19.1 | 18.0 | 1,822 | 10.2 |
| | 2001 | 616 | 5.4 | 15.4 | 2,583 | 3.5 | 3.0 | 18.0 | 1,249 | 19.3 | 18.4 | 1,777 | 8.7 |
| | 2000 | 579 | 4.1 | 15.5 | 2,541 | 3.8 | 2.0 | 19.1 | 1,244 | 21.7 | 20.3 | 1,860 | 8.9 |
| 1999 | 602 | 5.0 | 15.9 | 2,511 | 3.6 | 2.3 | 17.3 | 1,230 | 22.8 | 18.8 | 1,734 | 10.3 | |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 3
Run Date: 3 / 13 / 09

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

| Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | | | | | | | |
|--|------|----------------------------|--------------|-------------------------|-------------------------|--------------|------------------------------|------------|--------|-------------|------------|-----------------------------|--------|
| Race/Ethnicity and Year | | Birthweight ⁽⁴⁾ | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (7) | |
| | | Low | High | Short Stature (5) | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | | | |
| | | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th - | % <95th | % ≥95th | Number |
| Asian/Pacific Islander | 2008 | 1,337 | 7.6 | 3.3 | 5,549 | 10.4 | 2.8 | 14.5 | 2,595 | 17.2 | 16.2 | 4,119 | 9.2 |
| | 2007 | 1,293 | 7.0 | 3.3 | 5,258 | 10.5 | 2.8 | 13.5 | 2,398 | 17.3 | 15.6 | 3,825 | 9.6 |
| | 2006 | 1,240 | 6.8 | 4.2 | 5,340 | 11.8 | 3.2 | 13.8 | 2,531 | 16.0 | 15.1 | 4,073 | 10.8 |
| | 2005 | 1,232 | 7.5 | 3.7 | 5,312 | 12.3 | 3.8 | 13.0 | 2,544 | 16.9 | 15.8 | 3,743 | 13.1 |
| | 2004 | 1,279 | 7.3 | 4.7 | 5,209 | 11.8 | 3.6 | 13.5 | 2,537 | 17.1 | 15.8 | 3,710 | 13.7 |
| | 2003 | 1,143 | 6.7 | 4.5 | 4,803 | 9.2 | 3.4 | 15.2 | 2,409 | 17.8 | 18.3 | 3,410 | 13.3 |
| | 2002 | 1,057 | 8.4 | 5.0 | 4,877 | 9.9 | 3.5 | 15.4 | 2,420 | 17.4 | 18.4 | 3,595 | 13.8 |
| | 2001 | 1,159 | 6.1 | 4.5 | 5,159 | 9.3 | 3.7 | 14.7 | 2,646 | 17.6 | 18.2 | 3,701 | 11.5 |
| | 2000 | 1,162 | 6.7 | 5.8 | 5,416 | 9.6 | 2.7 | 15.8 | 2,931 | 16.9 | 18.4 | 4,024 | 13.1 |
| | 1999 | 1,064 | 8.0 | 4.7 | 5,699 | 9.7 | 3.0 | 14.3 | 3,272 | 18.4 | 15.9 | 4,343 | 12.6 |
| Multiple Races | 2008 | 1,613 | 9.1 | 8.4 | 6,639 | 4.5 | 3.3 | 14.6 | 2,994 | 16.8 | 13.0 | 4,895 | 10.2 |
| | 2007 | 1,512 | 9.0 | 8.6 | 6,121 | 4.6 | 3.3 | 13.1 | 2,628 | 17.3 | 12.3 | 4,352 | 10.7 |
| | 2006 | 1,652 | 7.7 | 7.4 | 6,053 | 5.2 | 3.7 | 13.2 | 2,473 | 16.1 | 13.1 | 4,289 | 10.0 |
| | 2005 | 376 | 10.4 | 8.2 | 1,076 | 5.4 | 4.3 | 14.2 | 343 | 14.3 | 12.5 | 723 | 9.3 |
| | 2004 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| | 2003 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| | 2002 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| | 2001 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| | 2000 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |
| | 1999 | 0 | * | * | 0 | * | * | * | 0 | * | * | 0 | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 18C
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 4
Run Date: 3 / 13 / 09

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

| Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | | | | | | | |
|--|------|----------------------------|--------------|-------------------------|-------------------------|--------------|------------------------------|------------|--------|-------------|-----------------------------|------------|--------|
| Race/Ethnicity and Year | | Birthweight ⁽⁴⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct (7) | | |
| | | Low | High | Short Stature (5) | Under- weight (6) | Obese (6) | ≥ 2 Yrs Overweight (6) | | | | | | |
| | | Number | % <2500 g | % >4000 g | Number | % <5th | % <5th | % ≥95th | Number | % 85th - | % <95th | % ≥95th | Number |
| All Other/Unknown | 2008 | 0 | * | * | 14 | * | * | * | 14 | * | * | 25 | * |
| | 2007 | 0 | * | * | 9 | * | * | * | 8 | * | * | 14 | * |
| | 2006 | 0 | * | * | 416 | 5.0 | 3.4 | 14.2 | 46 | * | * | 88 | * |
| | 2005 | 490 | 11.0 | 8.6 | 1,520 | 7.6 | 5.2 | 9.8 | 435 | 12.9 | 11.5 | 796 | 11.2 |
| | 2004 | 1 | * | * | 3 | * | * | * | 2 | * | * | 2 | * |
| | 2003 | 1 | * | * | 3 | * | * | * | 1 | * | * | 1 | * |
| | 2002 | 1 | * | * | 7 | * | * | * | 3 | * | * | 3 | * |
| | 2001 | 4 | * | * | 21 | * | * | * | 11 | * | * | 18 | * |
| | 2000 | 1 | * | * | 13 | * | * | * | 5 | * | * | 9 | * |
| | 1999 | 0 | * | * | 4 | * | * | * | 1 | * | * | 1 | * |
| Total All Race/Ethnic Groups | 2008 | 31,940 | 8.5 | 7.6 | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 | 11.0 |
| | 2007 | 31,162 | 8.7 | 7.5 | 120,071 | 5.4 | 3.7 | 12.2 | 51,636 | 16.1 | 13.1 | 84,006 | 11.3 |
| | 2006 | 31,039 | 8.6 | 7.6 | 118,548 | 5.7 | 3.8 | 12.4 | 51,667 | 16.3 | 13.0 | 85,045 | 10.4 |
| | 2005 | 28,451 | 8.6 | 8.1 | 112,753 | 5.5 | 4.5 | 12.3 | 51,410 | 16.0 | 13.0 | 77,037 | 12.2 |
| | 2004 | 29,547 | 8.7 | 8.3 | 113,275 | 5.4 | 4.3 | 12.4 | 51,825 | 16.3 | 13.3 | 78,255 | 13.2 |
| | 2003 | 28,471 | 8.3 | 8.5 | 109,384 | 5.2 | 4.4 | 12.2 | 50,284 | 15.9 | 13.0 | 75,448 | 12.2 |
| | 2002 | 27,201 | 8.5 | 8.7 | 106,429 | 5.5 | 4.3 | 11.8 | 48,993 | 15.0 | 11.8 | 74,420 | 12.7 |
| | 2001 | 27,782 | 8.4 | 8.9 | 104,796 | 5.6 | 4.6 | 11.4 | 47,741 | 14.6 | 11.4 | 71,952 | 13.0 |
| | 2000 | 27,000 | 8.5 | 8.9 | 102,004 | 5.5 | 4.5 | 11.2 | 46,744 | 15.1 | 11.5 | 70,955 | 13.6 |
| | 1999 | 25,565 | 8.8 | 9.0 | 100,387 | 5.4 | 4.4 | 10.5 | 47,584 | 14.6 | 10.1 | 69,517 | 13.5 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 19C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Summary of Trends in Breastfeeding Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | |
|--|------|------------------------------|------------------------|--------|--|--------|---|
| Race/Ethnicity and Year | | Breastfeeding ⁽⁴⁾ | | | | | |
| | | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| White, Not Hispanic | 2008 | 15,057 | 66.0 | 6,543 | 27.0 | 8,782 | 11.2 |
| | 2007 | 14,556 | 64.2 | 6,392 | 26.3 | 8,261 | 11.0 |
| | 2006 | 14,311 | 63.9 | 5,711 | 25.4 | 7,654 | 12.5 |
| | 2005 | 14,228 | 60.9 | 5,173 | 23.8 | 8,523 | 15.5 |
| | 2004 | 14,658 | 60.7 | 6,085 | 23.8 | 9,201 | 15.7 |
| | 2003 | 15,434 | 59.8 | 5,742 | 24.0 | 8,704 | 14.4 |
| | 2002 | 14,598 | 57.6 | 5,820 | 23.1 | 8,763 | 15.1 |
| | 2001 | 14,943 | 56.9 | 5,649 | 24.3 | 8,683 | 15.0 |
| | 2000 | 14,801 | 55.4 | 5,429 | 23.4 | 8,716 | 15.1 |
| | 1999 | 14,226 | 54.7 | 5,687 | 21.6 | 8,829 | 10.9 |
| Black, Not Hispanic | 2008 | 6,222 | 50.4 | 2,447 | 13.4 | 3,810 | 5.8 |
| | 2007 | 6,197 | 48.0 | 2,225 | 15.6 | 3,514 | 6.2 |
| | 2006 | 6,027 | 47.2 | 2,230 | 12.9 | 3,272 | 7.2 |
| | 2005 | 5,917 | 44.8 | 1,920 | 13.5 | 3,401 | 9.8 |
| | 2004 | 5,921 | 43.6 | 2,223 | 14.9 | 3,655 | 10.7 |
| | 2003 | 6,305 | 43.0 | 2,175 | 15.1 | 3,558 | 9.6 |
| | 2002 | 6,054 | 37.9 | 2,248 | 13.5 | 3,662 | 8.8 |
| | 2001 | 6,284 | 35.1 | 2,217 | 12.5 | 3,731 | 8.8 |
| | 2000 | 6,335 | 34.3 | 2,200 | 13.0 | 3,835 | 8.7 |
| | 1999 | 6,142 | 32.9 | 2,195 | 11.8 | 3,765 | 7.7 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 19C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 13 / 09

| Summary of Trends in Breastfeeding Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | |
|--|------|------------------------------|------------------------|--------|--|--------|---|
| Race/Ethnicity and Year | | Breastfeeding ⁽⁴⁾ | | | | | |
| | | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| Hispanic | 2008 | 7,334 | 78.6 | 3,503 | 40.2 | 5,002 | 21.0 |
| | 2007 | 7,124 | 78.7 | 3,575 | 37.8 | 4,953 | 20.6 |
| | 2006 | 7,274 | 76.8 | 3,289 | 39.0 | 4,374 | 20.6 |
| | 2005 | 5,483 | 76.8 | 2,201 | 36.5 | 3,780 | 24.3 |
| | 2004 | 4,837 | 77.5 | 2,166 | 38.0 | 3,518 | 22.6 |
| | 2003 | 4,831 | 76.3 | 1,970 | 35.0 | 3,303 | 21.8 |
| | 2002 | 4,712 | 72.3 | 1,935 | 32.8 | 3,159 | 21.7 |
| | 2001 | 4,646 | 70.6 | 1,773 | 32.9 | 2,794 | 20.8 |
| | 2000 | 3,984 | 67.0 | 1,535 | 31.3 | 2,426 | 20.2 |
| | 1999 | 3,346 | 65.1 | 1,322 | 27.8 | 2,157 | 16.4 |
| American Indian/Alaskan Native | 2008 | 425 | 58.8 | 221 | 21.3 | 316 | 11.7 |
| | 2007 | 493 | 55.8 | 244 | 20.5 | 391 | 9.2 |
| | 2006 | 570 | 62.1 | 232 | 24.1 | 376 | 12.8 |
| | 2005 | 643 | 60.3 | 223 | 24.7 | 427 | 16.2 |
| | 2004 | 604 | 59.6 | 234 | 28.2 | 438 | 18.3 |
| | 2003 | 586 | 59.4 | 252 | 21.8 | 440 | 19.8 |
| | 2002 | 583 | 60.0 | 249 | 30.1 | 418 | 19.9 |
| | 2001 | 594 | 54.5 | 214 | 20.1 | 403 | 18.4 |
| | 2000 | 556 | 52.3 | 255 | 23.5 | 433 | 22.2 |
| | 1999 | 587 | 55.7 | 213 | 25.4 | 375 | 11.2 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 19C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page: 3
Run Date: 3 / 13 / 09

| Summary of Trends in Breastfeeding Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | |
|--|------|------------------------------|------------------------|--------|--|--------|---|
| Race/Ethnicity and Year | | Breastfeeding ⁽⁴⁾ | | | | | |
| | | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| Asian/Pacific Islander | 2008 | 1,338 | 44.6 | 546 | 16.8 | 876 | 7.4 |
| | 2007 | 1,292 | 46.4 | 528 | 15.0 | 844 | 7.8 |
| | 2006 | 1,234 | 44.4 | 522 | 16.7 | 803 | 9.2 |
| | 2005 | 1,227 | 43.0 | 475 | 21.3 | 814 | 17.9 |
| | 2004 | 1,192 | 46.5 | 530 | 26.2 | 793 | 15.8 |
| | 2003 | 1,131 | 42.2 | 417 | 14.1 | 670 | 9.9 |
| | 2002 | 1,052 | 37.0 | 459 | 15.3 | 751 | 11.3 |
| | 2001 | 1,159 | 37.1 | 461 | 10.6 | 784 | 9.9 |
| | 2000 | 1,161 | 31.1 | 449 | 17.1 | 784 | 15.9 |
| | 1999 | 1,059 | 34.1 | 440 | 15.5 | 793 | 10.7 |
| Multiple Races | 2008 | 1,609 | 60.0 | 703 | 20.9 | 1,076 | 8.1 |
| | 2007 | 1,509 | 60.3 | 652 | 18.4 | 989 | 7.9 |
| | 2006 | 1,655 | 58.9 | 680 | 17.2 | 973 | 8.6 |
| | 2005 | 376 | 57.2 | 218 | 17.9 | 247 | 15.8 |
| | 2004 | 0 | * | 0 | * | 0 | * |
| | 2003 | 0 | * | 0 | * | 0 | * |
| | 2002 | 0 | * | 0 | * | 0 | * |
| | 2001 | 0 | * | 0 | * | 0 | * |
| | 2000 | 0 | * | 0 | * | 0 | * |
| | 1999 | 0 | * | 0 | * | 0 | * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 19C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Breastfeeding Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page: 4
Run Date: 3 / 13 / 09

| Summary of Trends in Breastfeeding Indicators by Race/Ethnicity ⁽³⁾ | | | | | | | |
|--|------|------------------------------|------------------------|--------|--|--------|---|
| Race/Ethnicity and Year | | Breastfeeding ⁽⁴⁾ | | | | | |
| | | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months |
| All Other/Unknown | 2008 | 0 | * | 0 | * | 0 | * |
| | 2007 | 0 | * | 0 | * | 0 | * |
| | 2006 | 0 | * | 62 | * | 15 | * |
| | 2005 | 736 | 64.7 | 199 | 16.6 | 225 | 12.9 |
| | 2004 | 1 | * | 0 | * | 0 | * |
| | 2003 | 1 | * | 0 | * | 0 | * |
| | 2002 | 1 | * | 0 | * | 1 | * |
| | 2001 | 4 | * | 0 | * | 1 | * |
| | 2000 | 1 | * | 0 | * | 0 | * |
| | 1999 | 0 | * | 0 | * | 0 | * |
| Total All Race/Ethnic Groups | 2008 | 31,985 | 64.6 | 13,963 | 27.1 | 19,862 | 12.3 |
| | 2007 | 31,171 | 63.2 | 13,616 | 26.6 | 18,952 | 12.3 |
| | 2006 | 31,071 | 62.6 | 12,726 | 26.0 | 17,467 | 13.2 |
| | 2005 | 28,610 | 59.9 | 10,409 | 24.2 | 17,417 | 16.4 |
| | 2004 | 27,213 | 59.3 | 11,238 | 25.0 | 17,605 | 16.1 |
| | 2003 | 28,288 | 58.2 | 10,556 | 23.8 | 16,675 | 14.8 |
| | 2002 | 27,000 | 55.0 | 10,711 | 22.6 | 16,754 | 14.9 |
| | 2001 | 27,630 | 53.4 | 10,314 | 22.5 | 16,396 | 14.4 |
| | 2000 | 26,838 | 51.0 | 9,868 | 22.1 | 16,194 | 14.6 |
| | 1999 | 25,360 | 49.9 | 9,857 | 20.0 | 15,919 | 10.9 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 20C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Summary of Trends in Growth Indicators by Age ⁽³⁾ | | | | | | | | | | | | | | | | | | |
|--|--------|-------------------|------|------------------|------------|------------------------|---|-------|------|----------------|------|-------------------|-------|------------------|-----------|------------------------|-------|---|
| Age and Year * | | Height and Weight | | | | | | | | Age and Year * | | Height and Weight | | | | | | |
| | | Short Stature (4) | | Under-weight (5) | Obese (5) | ≥ 2 Yrs Overweight (5) | | | | | | Short Stature (4) | | Under-weight (5) | Obese (5) | ≥ 2 Yrs Overweight (5) | | |
| | | Number | % | % | % | Number | % | % | | | | Number | % | % | % | Number | % | % |
| | | <5th | <5th | ≥95th | 85th-<95th | | | ≥95th | | | <5th | <5th | ≥95th | 85th-<95th | | | ≥95th | |
| 0 - 11 Months | | | | | | | | | | 24 - 35 Months | | | | | | | | |
| 2008 | 43,214 | 7.5 | 4.9 | 10.5 | 0 | * | * | | 2008 | 21,266 | 4.2 | 2.9 | 12.2 | 21,266 | 15.7 | 12.2 | | |
| 2007 | 42,270 | 7.2 | 5.3 | 9.2 | 0 | * | * | | 2007 | 19,319 | 4.2 | 3.2 | 11.7 | 19,319 | 15.6 | 11.7 | | |
| 2006 | 42,479 | 7.5 | 5.5 | 9.7 | 0 | * | * | | 2006 | 19,223 | 4.4 | 3.0 | 11.7 | 19,223 | 16.0 | 11.7 | | |
| 2005 | 34,813 | 7.4 | 8.0 | 8.4 | 0 | * | * | | 2005 | 18,912 | 4.3 | 3.4 | 11.5 | 18,912 | 15.6 | 11.5 | | |
| 2004 | 35,031 | 7.6 | 7.4 | 8.1 | 0 | * | * | | 2004 | 18,993 | 4.1 | 3.2 | 11.3 | 18,993 | 15.8 | 11.3 | | |
| 2003 | 34,087 | 7.3 | 6.9 | 8.2 | 0 | * | * | | 2003 | 18,447 | 3.7 | 3.7 | 11.5 | 18,447 | 15.5 | 11.5 | | |
| 2002 | 32,759 | 7.6 | 6.4 | 8.8 | 0 | * | * | | 2002 | 18,003 | 3.8 | 4.2 | 10.5 | 18,003 | 13.8 | 10.5 | | |
| 2001 | 32,739 | 8.1 | 6.6 | 8.9 | 0 | * | * | | 2001 | 17,642 | 3.5 | 4.5 | 9.8 | 17,642 | 13.4 | 9.8 | | |
| 2000 | 31,190 | 7.5 | 7.2 | 7.8 | 0 | * | * | | 2000 | 17,219 | 3.7 | 4.0 | 9.8 | 17,219 | 13.9 | 9.8 | | |
| 1999 | 29,268 | 7.7 | 6.8 | 7.8 | 0 | * | * | | 1999 | 17,215 | 3.4 | 4.1 | 9.2 | 17,215 | 12.8 | 9.2 | | |
| 12 - 23 Months | | | | | | | | | | 36 - 47 Months | | | | | | | | |
| 2008 | 27,680 | 5.6 | 2.0 | 16.9 | 0 | * | * | | 2008 | 17,663 | 3.9 | 3.2 | 13.8 | 17,663 | 16.5 | 13.8 | | |
| 2007 | 26,165 | 5.4 | 2.1 | 15.4 | 0 | * | * | | 2007 | 16,798 | 4.0 | 3.3 | 13.8 | 16,798 | 16.2 | 13.8 | | |
| 2006 | 24,402 | 5.7 | 2.1 | 15.5 | 0 | * | * | | 2006 | 16,557 | 4.4 | 3.3 | 13.2 | 16,557 | 16.5 | 13.2 | | |
| 2005 | 26,530 | 5.9 | 2.1 | 16.0 | 0 | * | * | | 2005 | 16,587 | 4.1 | 3.4 | 13.5 | 16,587 | 15.7 | 13.5 | | |
| 2004 | 26,419 | 5.4 | 2.2 | 16.1 | 0 | * | * | | 2004 | 16,748 | 3.9 | 3.5 | 14.3 | 16,748 | 16.1 | 14.3 | | |
| 2003 | 25,013 | 5.4 | 2.4 | 16.0 | 0 | * | * | | 2003 | 16,353 | 3.9 | 4.0 | 13.6 | 16,353 | 15.7 | 13.6 | | |
| 2002 | 24,677 | 5.9 | 2.4 | 15.6 | 0 | * | * | | 2002 | 15,852 | 4.1 | 4.2 | 12.2 | 15,852 | 15.1 | 12.2 | | |
| 2001 | 24,316 | 5.8 | 2.5 | 14.9 | 0 | * | * | | 2001 | 15,447 | 3.8 | 4.5 | 11.6 | 15,447 | 14.8 | 11.6 | | |
| 2000 | 24,070 | 5.6 | 2.4 | 15.1 | 0 | * | * | | 2000 | 14,940 | 4.5 | 3.6 | 12.4 | 14,940 | 15.0 | 12.4 | | |
| 1999 | 23,535 | 5.5 | 2.5 | 14.6 | 0 | * | * | | 1999 | 15,166 | 4.3 | 4.0 | 10.7 | 15,166 | 15.4 | 10.7 | | |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 20C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Growth Indicators by Age
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 13 / 09

| Summary of Trends in Growth Indicators by Age ⁽³⁾ | | | | | | | | | | | | | | | | | | | |
|--|--|-------------------|------|-------------------|------------|------------------------|------|-------|--|--|----------------|------|-------------------|------------|-------------------|-----------|------------------------|---|---|
| Age and Year * | | Height and Weight | | | | | | | | | Age and Year * | | Height and Weight | | | | | | |
| | | Short Stature (4) | | Under- weight (5) | Obese (5) | ≥ 2 Yrs Overweight (5) | | | | | | | Short Stature (4) | | Under- weight (5) | Obese (5) | ≥ 2 Yrs Overweight (5) | | |
| | | Number | % | % | % | Number | % | % | | | | | Number | % | % | % | Number | % | % |
| | | <5th | <5th | ≥95th | 85th-<95th | | | ≥95th | | | <5th | <5th | ≥95th | 85th-<95th | | | ≥95th | | |
| 48 - 59 Months | | | | | | | | | | | | | | | | | | | |
| 2008 | | 16,946 | 3.6 | 3.0 | 15.1 | 16,946 | 16.7 | 15.1 | | | | | | | | | | | |
| 2007 | | 15,519 | 3.8 | 2.9 | 14.2 | 15,519 | 16.5 | 14.2 | | | | | | | | | | | |
| 2006 | | 15,887 | 3.8 | 3.0 | 14.5 | 15,887 | 16.7 | 14.5 | | | | | | | | | | | |
| 2005 | | 15,911 | 3.7 | 3.3 | 14.2 | 15,911 | 16.9 | 14.2 | | | | | | | | | | | |
| 2004 | | 16,084 | 3.8 | 3.3 | 14.7 | 16,084 | 17.2 | 14.7 | | | | | | | | | | | |
| 2003 | | 15,484 | 3.6 | 3.3 | 14.3 | 15,484 | 16.6 | 14.3 | | | | | | | | | | | |
| 2002 | | 15,138 | 3.7 | 3.4 | 13.0 | 15,138 | 16.4 | 13.0 | | | | | | | | | | | |
| 2001 | | 14,652 | 3.8 | 3.9 | 13.0 | 14,652 | 16.0 | 13.0 | | | | | | | | | | | |
| 2000 | | 14,585 | 4.2 | 3.6 | 12.5 | 14,585 | 16.7 | 12.5 | | | | | | | | | | | |
| 1999 | | 15,203 | 4.4 | 3.6 | 10.7 | 15,203 | 15.9 | 10.7 | | | | | | | | | | | |
| Total All Age Groups | | | | | | | | | | | | | | | | | | | |
| 2008 | | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | | | | | | | | | | | |
| 2007 | | 120,071 | 5.4 | 3.7 | 12.2 | 51,636 | 16.1 | 13.1 | | | | | | | | | | | |
| 2006 | | 118,548 | 5.7 | 3.8 | 12.4 | 51,667 | 16.3 | 13.0 | | | | | | | | | | | |
| 2005 | | 112,753 | 5.5 | 4.5 | 12.3 | 51,410 | 16.0 | 13.0 | | | | | | | | | | | |
| 2004 | | 113,275 | 5.4 | 4.3 | 12.4 | 51,825 | 16.3 | 13.3 | | | | | | | | | | | |
| 2003 | | 109,384 | 5.2 | 4.4 | 12.2 | 50,284 | 15.9 | 13.0 | | | | | | | | | | | |
| 2002 | | 106,429 | 5.5 | 4.3 | 11.8 | 48,993 | 15.0 | 11.8 | | | | | | | | | | | |
| 2001 | | 104,796 | 5.6 | 4.6 | 11.4 | 47,741 | 14.6 | 11.4 | | | | | | | | | | | |
| 2000 | | 102,004 | 5.5 | 4.5 | 11.2 | 46,744 | 15.1 | 11.5 | | | | | | | | | | | |
| 1999 | | 100,387 | 5.4 | 4.4 | 10.5 | 47,584 | 14.6 | 10.1 | | | | | | | | | | | |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(5) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 21C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 13 / 09

| Summary of Trends in Anemia by Age (3) | | | | | | |
|--|------|--------------------------|------|----------------|------|-------------|
| Age and Year | | Anemia Low Hb/Hct (4) | | | | |
| | | Number | % | | | |
| 6 - 11 Months | | | | 18 - 23 Months | | |
| | 2008 | 5,410 | 16.2 | | 2008 | 14,367 12.5 |
| | 2007 | 3,416 | 15.8 | | 2007 | 13,253 12.8 |
| | 2006 | 6,956 | 16.2 | | 2006 | 12,474 12.1 |
| | 2005 | 5,354 | 16.1 | | 2005 | 11,613 13.8 |
| | 2004 | 5,135 | 16.9 | | 2004 | 11,660 15.1 |
| | 2003 | 5,357 | 16.1 | | 2003 | 11,332 14.3 |
| | 2002 | 5,411 | 15.4 | | 2002 | 11,117 15.0 |
| | 2001 | 4,964 | 16.5 | | 2001 | 10,697 15.6 |
| | 2000 | 4,330 | 19.0 | | 2000 | 10,306 16.5 |
| | 1999 | 4,340 | 17.7 | | 1999 | 9,909 15.7 |
| 12 - 17 Months | | | | 24 - 35 Months | | |
| | 2008 | 18,196 | 16.8 | | 2008 | 21,905 10.2 |
| | 2007 | 17,937 | 16.8 | | 2007 | 19,796 11.0 |
| | 2006 | 16,217 | 15.5 | | 2006 | 19,862 9.8 |
| | 2005 | 15,972 | 16.1 | | 2005 | 16,657 12.5 |
| | 2004 | 16,255 | 17.2 | | 2004 | 16,750 13.8 |
| | 2003 | 15,184 | 15.4 | | 2003 | 16,205 12.7 |
| | 2002 | 15,081 | 15.6 | | 2002 | 16,010 13.9 |
| | 2001 | 14,814 | 16.2 | | 2001 | 15,688 13.8 |
| | 2000 | 14,755 | 17.9 | | 2000 | 15,665 14.0 |
| | 1999 | 14,272 | 16.8 | | 1999 | 15,384 14.6 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 21C
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Anemia by Age
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 13 / 09

| Summary of Trends in Anemia by Age (3) | | | | | | | | |
|--|------|--------------------------|-----|--|----------------------|------|--------------------------|------|
| Age and Year | | Anemia Low Hb/Hct (4) | | | Age and Year | | Anemia Low Hb/Hct (4) | |
| | | Number | % | | | | Number | % |
| 36 - 59 Months | | | | | Total All Age Groups | | | |
| | 2008 | 31,741 | 6.6 | | | 2008 | 91,619 | 11.0 |
| | 2007 | 29,604 | 6.9 | | | 2007 | 84,006 | 11.3 |
| | 2006 | 29,536 | 6.1 | | | 2006 | 85,045 | 10.4 |
| | 2005 | 27,441 | 8.2 | | | 2005 | 77,037 | 12.2 |
| | 2004 | 28,455 | 9.2 | | | 2004 | 78,255 | 13.2 |
| | 2003 | 27,370 | 8.4 | | | 2003 | 75,448 | 12.2 |
| | 2002 | 26,801 | 8.9 | | | 2002 | 74,420 | 12.7 |
| | 2001 | 25,789 | 9.1 | | | 2001 | 71,952 | 13.0 |
| | 2000 | 25,899 | 9.0 | | | 2000 | 70,955 | 13.6 |
| | 1999 | 25,612 | 9.5 | | | 1999 | 69,517 | 13.5 |
| All Other/Unknown | | | | | | | | |
| | 2008 | 0 | * | | | | | |
| | 2007 | 0 | * | | | | | |
| | 2006 | 0 | * | | | | | |
| | 2005 | 0 | * | | | | | |
| | 2004 | 0 | * | | | | | |
| | 2003 | 0 | * | | | | | |
| | 2002 | 0 | * | | | | | |
| | 2001 | 0 | * | | | | | |
| | 2000 | 0 | * | | | | | |
| | 1999 | 0 | * | | | | | |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

